Anorectal manometry and Hirschsprung's disease

Access this article online	
Quick Response Code:	Website: www.afrjpaedsurg.org
	DOI: 10.4103/0189-6725.120885

Sir,

The recent report on anorectal manometry in following up post-operative Hirschsprung's disease is very interesting.[1] Demirbag *et al.*, concluded that "the majority of the patients have impaired anorectal motility.[1] Indeed, Hirschsprung's disease is a congenital disease which can be confirmed by histopathological defect. The surgical management of this disease is usually for supporting and correction. The resolution of the anorectal function is hardly to be resolved. The use of anorectal manometry might be limitedly useful in post-operative following up of the patients with Hirschsprung's disease. However, the use of anorectal manometry might be useful for classification of the severity of dysganglionosis for help reach diagnosis in the suspicious patients with Hirschsprung's disease.[2,3]

Somsri Wiwanitkit, Viroj Wiwanitkit[1]

Wiwanitkit House, Bangkhae, Bangkok, Thailand, [1]Joseph Ayobabalola University, Ikeji-Arakeji, Osun State, Nigeria
E-mail: somsriwiwan@hotmail.com

REFERENCES

1. Demirbag S, Tiryaki T, Purtuloglu T. Importance of anorectal manometry after definitive surgery for Hirschsprung's disease in children. Afr J Paediatr Surg 2013;10:1-4.
2. Wu XJ, Zhang HY, Li N, Yan MS, Wei J, Yu DH, *et al*. A new diagnostic scoring system to differentiate Hirschsprung's disease from Hirschsprung's disease-allied disorders in patients with suspected intestinal dysganglionosis. Int J Colorectal Dis 2013.
3. Zhou XL, Chen FB, Ou BY, Zhang XP, Jiang MZ. Evaluation of clinical value of anorectal manometry for diagnosis of Hirschsprung's disease in neonate. Zhonghua Er Ke Za Zhi 2004;42:681-3.